

A SAT Solver's Opinion on the Erdős-Faber-Lovász Conjecture

Markus Kirchweger


Tomáš Peitl

Stefan Szeider

SAT 2023

Jul 6th


Outline




In this talk

What conjecture


How we tackle it





EFL Conjecture ('72)

Every **linear** hypergraph with n vertices is n -edge-colorable.



EFL Conjecture ('72)

Every **linear** hypergraph with n vertices is n -edge-colorable.

Theorem ([KKK⁺21])

There is n_0 s.t. when $n > n_0$, every linear hypergraph with n vertices is n -edge-colorable.

$F_n(X) \Leftrightarrow X$ is a linear hypergraph with n vertices not edge-colorable with n colors

$F_n(X) \Leftrightarrow X$ is a linear hypergraph with n vertices not edge-colorable with n colors

$F_n(X) \rightarrow \sim\text{SAT solver}$

$F_n(X) \Leftrightarrow X$ is a linear hypergraph with n vertices not edge-colorable with n colors

$F_n(X) \rightarrow \sim\text{SAT solver}$

- we will try to generate a counterexample to EFL with [SAT Modulo Symmetries \(SMS\)](#) [KS21];

$F_n(X) \Leftrightarrow X$ is a linear hypergraph with n vertices not edge-colorable with n colors

$F_n(X) \rightarrow \sim\text{SAT solver}$

- we will try to generate a counterexample to EFL with [SAT Modulo Symmetries \(SMS\)](#) [KS21];
- previously, the conjecture has been verified for $n \leq 12$ [Hin81, RAP14]; we verify and extend this;

$F_n(X) \Leftrightarrow X$ is a linear hypergraph with n vertices not edge-colorable with n colors

$F_n(X) \rightarrow \sim\text{SAT solver}$


- we will try to generate a counterexample to EFL with **SAT Modulo Symmetries (SMS)** [KS21];
- previously, the conjecture has been verified for $n \leq 12$ [Hin81, RAP14]; we verify and extend this;
- we also verify a **generalized** conjecture and generate **extremal hypergraphs** (which need n colors);

$F_n(X) \Leftrightarrow X$ is a linear hypergraph with n vertices not edge-colorable with n colors

$F_n(X) \rightarrow \sim\text{SAT solver}$


- we will try to generate a counterexample to EFL with [SAT Modulo Symmetries \(SMS\)](#) [KS21];
- previously, the conjecture has been verified for $n \leq 12$ [Hin81, RAP14]; we verify and extend this;
- we also verify a [generalized](#) conjecture and generate [extremal hypergraphs](#) (which need n colors);
- inspired by the generated examples, we prove some [general theorems](#) about (non-)extremality;

Overview of “~SAT solver”




- **Generator** is a SAT solver that enumerates models modulo symmetries;

Overview of “~SAT solver”




- **Generator** is a SAT solver that enumerates models modulo symmetries;
- **Solution candidate** is a linear hypergraph that passes F_n , but can be colorable;

Overview of “~SAT solver”



- **Generator** is a SAT solver that enumerates models modulo symmetries;
- **Solution candidate** is a linear hypergraph that passes F_n , but can be colorable;
- non-colorability cannot be encoded in polynomial size, so we use a **2nd SAT solver**;

Overview of “~SAT solver”



- **Generator** is a SAT solver that enumerates models modulo symmetries;
- **Solution candidate** is a linear hypergraph that passes F_n , but can be colorable;
- non-colorability cannot be encoded in polynomial size, so we use a **2nd SAT solver**;
- SMS relies on a custom propagator (in Cadical), the coloring is “incremental”.

SAT Modulo Symmetries for Hypergraphs


- SMS was recently proposed for **graph** generation modulo isomorphisms (symmetries);

SAT Modulo Symmetries for Hypergraphs

- SMS was recently proposed for **graph** generation modulo isomorphisms (symmetries);
- avoids the bottlenecks of large static symmetry breaking and generate-and-test methods;


SAT Modulo Symmetries for Hypergraphs

- SMS was recently proposed for **graph** generation modulo isomorphisms (symmetries);
- avoids the bottlenecks of large static symmetry breaking and generate-and-test methods;
- works with graphs, but we can express hypergraphs as **incidence graphs**;



SAT Modulo Symmetries for Hypergraphs


- SMS was recently proposed for **graph** generation modulo isomorphisms (symmetries);
- avoids the bottlenecks of large static symmetry breaking and generate-and-test methods;
- works with graphs, but we can express hypergraphs as **incidence graphs**;



- pretend that edges between vertices and between hyperedges are also allowed, and say they are not there (with unit clauses);

SAT Modulo Symmetries for Hypergraphs

- SMS was recently proposed for **graph** generation modulo isomorphisms (symmetries);
- avoids the bottlenecks of large static symmetry breaking and generate-and-test methods;
- works with graphs, but we can express hypergraphs as **incidence graphs**;



- pretend that edges between vertices and between hyperedges are also allowed, and say they are not there (with unit clauses);
- linearity = no 4-cycle in the incidence graph.

Generating non- n -edge-colorable hypergraphs

- we can now express that we want a linear hypergraph with n vertices and m hyperedges with a formula of size $\text{poly}(n, m)$; but

Generating non- n -edge-colorable hypergraphs

- we can now express that we want a linear hypergraph with n vertices and m hyperedges with a formula of size $\text{poly}(n, m)$; but
- non-colorability requires exponential size;

Generating non- n -edge-colorable hypergraphs

- we can now express that we want a linear hypergraph with n vertices and m hyperedges with a formula of size $\text{poly}(n, m)$; but
- non-colorability requires exponential size;
- instead, we generate hypergraphs and test for colorability, and ...

Generating non- n -edge-colorable hypergraphs

- we can now express that we want a linear hypergraph with n vertices and m hyperedges with a formula of size $\text{poly}(n, m)$; but
- non-colorability requires exponential size;
- instead, we generate hypergraphs and test for colorability, and ...
- when a coloring $c : H \rightarrow [\chi]$ is found for a candidate, learn a clause that forbids this coloring;

$$\bigvee_{c(e_1)=c(e_2)} \mathcal{I}_{e_1, e_2}$$

Generating non- n -edge-colorable hypergraphs

- we can now express that we want a linear hypergraph with n vertices and m hyperedges with a formula of size $\text{poly}(n, m)$; but
- non-colorability requires exponential size;
- instead, we generate hypergraphs and test for colorability, and ...
- when a coloring $c : H \rightarrow [\chi]$ is found for a candidate, learn a clause that forbids this coloring;

$$\bigvee_{c(e_1)=c(e_2)} \mathcal{I}_{e_1, e_2}$$

- follows the co-certificate learning method [KPS23];

Generating non- n -edge-colorable hypergraphs

- we can now express that we want a linear hypergraph with n vertices and m hyperedges with a formula of size $\text{poly}(n, m)$; but
- non-colorability requires exponential size;
- instead, we generate hypergraphs and test for colorability, and ...
- when a coloring $c : H \rightarrow [\chi]$ is found for a candidate, learn a clause that forbids this coloring;

$$\bigvee_{c(e_1)=c(e_2)} \mathcal{I}_{e_1, e_2}$$

- follows the co-certificate learning method [KPS23];
- can also generate non- $(n - 1)$ -colorable (extremal) hypergraphs in the same manner;

Restrictions and generalizations

- adding edges is safe, so we can restrict to **linear spaces**;

Restrictions and generalizations

- adding edges is safe, so we can restrict to **linear spaces**;
- works for **both** counterexample search and extremal search;

Restrictions and generalizations

- adding edges is safe, so we can restrict to **linear spaces**;
- works for **both** counterexample search and extremal search;
- in graphs, the only linear space is the complete graph K_n ;

$$\chi'(K_n) = \begin{cases} n & n \text{ odd} \\ n - 1 & n \text{ even} \end{cases}$$

Restrictions and generalizations

- adding edges is safe, so we can restrict to **linear spaces**;
- works for **both** counterexample search and extremal search;
- in graphs, the only linear space is the complete graph K_n ;

$$\chi'(K_n) = \begin{cases} n & n \text{ odd} \\ n - 1 & n \text{ even} \end{cases}$$

- Vizing's Theorem: $\chi'(G) \in \{\Delta(G), \Delta(G) + 1\}$ ($\Delta = \max \text{ degree}$);

Restrictions and generalizations

- adding edges is safe, so we can restrict to **linear spaces**;
- works for **both** counterexample search and extremal search;
- in graphs, the only linear space is the complete graph K_n ;

$$\chi'(K_n) = \begin{cases} n & n \text{ odd} \\ n - 1 & n \text{ even} \end{cases}$$

- Vizing's Theorem: $\chi'(G) \in \{\Delta(G), \Delta(G) + 1\}$ ($\Delta = \max$ degree);
- does **not** work for linear hypergraphs, but a conjecture says:

Restrictions and generalizations

- adding edges is safe, so we can restrict to **linear spaces**;
- works for **both** counterexample search and extremal search;
- in graphs, the only linear space is the complete graph K_n ;

$$\chi'(K_n) = \begin{cases} n & n \text{ odd} \\ n - 1 & n \text{ even} \end{cases}$$

- Vizing's Theorem: $\chi'(G) \in \{\Delta(G), \Delta(G) + 1\}$ ($\Delta = \max \text{ degree}$);
- does **not** work for linear hypergraphs, but a conjecture says:

Conjecture (Füredi [Für86], Berge [Ber89])

Let $H = (V, E)$ be a linear hypergraph, and let $\zeta = \max_{v \in V} |\bigcup_{v \in e} e|$. Then H is ζ -edge-colorable. (For graphs $\zeta = \Delta + 1$.)


Results

Theorem

The EFL' Conjecture holds for n, m if $n \leq 12$; or

- *$n = 13$ and $m \in [13, 32] \cup [55, 78]$; or*
- *$n = 14$ and $m \in [14, 28] \cup [70, 91]$; or*
- *$n = 15$ and $m \in [15, 29] \cup [84, 105]$; or*
- *$n = 16$ and $m \in [16, 30] \cup \{99\} \cup [101, 120]$; or*
- *$n = 17$ and $m \in [17, 30] \cup [117, 136]$; or*
- *$n = 18$ and $m \in [18, 31] \cup [134, 153]$.*

Coloring time vs total time for $n = 12$



Coloring time vs total time for $n = 15$




Figure: The distribution of the running times for $n = 12$ and $n = 15$.

Theorem

The FB' Conjecture holds for n and m if $n \leq 10$; or

- *$n = 11$ and $m \in [11, 20] \cup [37, 55]$; or*
- *$n = 12$ and $m \in [12, 17] \cup [49, 66]$.*




Figure: The distribution of the running times for the EFL' and FB' conjectures for $n \in [9, 10]$.

Theorem

The linear spaces with $n \leq 12$ vertices and chromatic index n are precisely $\mathcal{H}_{n,k}$ for all $k \not\equiv n \pmod 2$, and additionally $\mathcal{H}_{7,3}$, $\mathcal{H}_{9,3}$, $\mathcal{H}_{11,3}$, $\mathcal{H}_{7,3 \setminus 3}$, $\mathcal{H}_{11,3 \cap 3}$, $\mathcal{H}_{11,3 \setminus 3}$, and the Fano plane.

K_n is extremal precisely when n is odd


n	5	6	7	8	9	10	11
	✓	✗	✓	✗	✓	✗	✓

$\mathcal{H}_{n,2}$ is extremal precisely when n is odd

n	5	6	7	8	9	10	11
k							
2	✓	✗	✓	✗	✓	✗	✓

$\mathcal{H}_{n,k}$ is the linear space with one k -edge and all the rest 2-edges.

$\mathcal{H}_{7,3} =$




$\mathcal{H}_{n,2}$ is extremal precisely when n is odd

n	5	6	7	8	9	10	11
k							
2	✓	✗	✓	✗	✓	✗	✓
3	✗	✓	✗	✓	✗	✓	✗

$\mathcal{H}_{n,k}$ is the linear space with one k -edge and all the rest 2-edges.

$\mathcal{H}_{7,3} =$




$\mathcal{H}_{n,2}$ is extremal precisely when n is odd

n	5	6	7	8	9	10	11
k							
2	✓	✗	✓	✗	✓	✗	✓
3	✗	✓	✗	✓	✗	✓	✗
4	✓	✗	✓	✗	✓	✗	✓

$\mathcal{H}_{n,k}$ is the linear space with one k -edge and all the rest 2-edges.

$\mathcal{H}_{7,3} =$




$\mathcal{H}_{n,2}$ is extremal precisely when n is odd

n	5	6	7	8	9	10	11
k							
2	✓	✗	✓	✗	✓	✗	✓
3	✗	✓	✗	✓	✗	✓	✗
4	✓	✗	✓	✗	✓	✗	✓
5		✓	✗	✓	✗	✓	✗

$\mathcal{H}_{n,k}$ is the linear space with one k -edge and all the rest 2-edges.

$\mathcal{H}_{7,3} =$




$\mathcal{H}_{n,2}$ is extremal precisely when n is odd

n	5	6	7	8	9	10	11
k							
2	✓	✗	✓	✗	✓	✗	✓
3	✗	✓	✗	✓	✗	✓	✗
4	✓	✗	✓	✗	✓	✗	✓
5		✓	✗	✓	✗	✓	✗
6			✓	✗	✓	✗	✓

$\mathcal{H}_{n,k}$ is the linear space with one k -edge and all the rest 2-edges.

$\mathcal{H}_{7,3} =$




$\mathcal{H}_{n,2}$ is extremal precisely when n is odd

n	5	6	7	8	9	10	11
k							
2	✓	✗	✓	✗	✓	✗	✓
3	✗	✓	✗	✓	✗	✓	✗
4	✓	✗	✓	✗	✓	✗	✓
5		✓	✗	✓	✗	✓	✗
6			✓	✗	✓	✗	✓
7				✓	✗	✓	✗

$\mathcal{H}_{n,k}$ is the linear space with one k -edge and all the rest 2-edges.

$\mathcal{H}_{7,3} =$




$\mathcal{H}_{n,2}$ is extremal precisely when n is odd

n	5	6	7	8	9	10	11
k							
2	✓	✗	✓	✗	✓	✗	✓
3	✗	✓	✗	✓	✗	✓	✗
4	✓	✗	✓	✗	✓	✗	✓
5		✓	✗	✓	✗	✓	✗
6			✓	✗	✓	✗	✓
7				✓	✗	✓	✗
8					✓	✗	✓

$\mathcal{H}_{n,k}$ is the linear space with one k -edge and all the rest 2-edges.

$\mathcal{H}_{7,3} =$




$\mathcal{H}_{n,2}$ is extremal precisely when n is odd

n	5	6	7	8	9	10	11
k							
2	✓	✗	✓	✗	✓	✗	✓
3	✗	✓	✗	✓	✗	✓	✗
4	✓	✗	✓	✗	✓	✗	✓
5		✓	✗	✓	✗	✓	✗
6			✓	✗	✓	✗	✓
7				✓	✗	✓	✗
8					✓	✗	✓
9						✓	✗

$\mathcal{H}_{n,k}$ is the linear space with one k -edge and all the rest 2-edges.

$\mathcal{H}_{7,3} =$




$\mathcal{H}_{n,2}$ is extremal precisely when n is odd

n	5	6	7	8	9	10	11
k							
2	✓	✗	✓	✗	✓	✗	✓
3	✗	✓	✗	✓	✗	✓	✗
4	✓	✗	✓	✗	✓	✗	✓
5		✓	✗	✓	✗	✓	✗
6			✓	✗	✓	✗	✓
7				✓	✗	✓	✗
8					✓	✗	✓
9						✓	✗
10							✓

$\mathcal{H}_{n,k}$ is the linear space with one k -edge and all the rest 2-edges.

$\mathcal{H}_{7,3} =$




Chain

$\mathcal{H}_{n,k}$ is extremal when $n \not\equiv k \pmod 2$, and

n	5	6	7	8	9	10	11
k							
2	✓	✗	✓	✗	✓	✗	✓
3	✗	✓	✗	✓	✗	✓	✗
4	✓	✗	✓	✗	✓	✗	✓
5		✓	✗	✓	✗	✓	✗
6			✓	✗	✓	✗	✓
7				✓	✗	✓	✗
8					✓	✗	✓
9						✓	✗
10							✓

$\mathcal{H}_{n,k}$ is the linear space with one k -edge and all the rest 2-edges.

$\mathcal{H}_{7,3} =$




Chain

$\mathcal{H}_{n,k}$ is extremal when $n \not\equiv k \pmod 2$, and sometimes when $n \equiv k \equiv 1 \pmod 2$

n	5	6	7	8	9	10	11
k							
2	✓	✗	✓	✗	✓	✗	✓
3	✗	✓	✓	✓	✓	✓	✓
4	✓	✗	✓	✗	✓	✗	✓
5		✓	✗	✓	✗	✓	✗
6			✓	✗	✓	✗	✓
7				✓	✗	✓	✗
8					✓	✗	✓
9						✓	✗
10							✓

$\mathcal{H}_{n,k}$ is the linear space with one k -edge and all the rest 2-edges.

$\mathcal{H}_{7,3} =$




Chain

$\mathcal{H}_{n,k}$ is extremal when $n \not\equiv k \pmod 2$, and sometimes when $n \equiv k \equiv 1 \pmod 2$, but never when $n \equiv k \equiv 0 \pmod 2$.

n	5	6	7	8	9	10	11
k							
2	✓	✗	✓	✗	✓	✗	✓
3	✗	✓	✓	✓	✓	✓	✓
4	✓	✗	✓	✗	✓	✗	✓
5		✓	✗	✓	✗	✓	✗
6			✓	✗	✓	✗	✓
7				✓	✗	✓	✗
8					✓	✗	✓
9						✓	✗
10							✓



$\mathcal{H}_{n,k}$ is the linear space with one k -edge and all the rest 2-edges.

$\mathcal{H}_{7,3} =$






More nice pictures

$\mathcal{H}_{7,3,7/3}$



More nice pictures



- SMS is great for (hyper)graph search with constraints, even for problems with a coNP-complete constraint; try it out:
 - ▶ <https://sat-modulo-symmetries.readthedocs.io>
 - ▶ <https://github.com/markirch/sat-modulo-symmetries>

Conclusion

- **SMS** is great for (hyper)graph search with constraints, even for problems with a coNP-complete constraint; try it out:
 - ▶ <https://sat-modulo-symmetries.readthedocs.io>
 - ▶ <https://github.com/markirch/sat-modulo-symmetries>
- a **SAT-based** exploration of small cases can inspire **general** results;

- SMS is great for (hyper)graph search with constraints, even for problems with a coNP-complete constraint; try it out:
 - ▶ <https://sat-modulo-symmetries.readthedocs.io>
 - ▶ <https://github.com/markirch/sat-modulo-symmetries>
- a SAT-based exploration of small cases can inspire general results;
- highly symmetric hypergraphs are quite hard for SMS;

- SMS is great for (hyper)graph search with constraints, even for problems with a coNP-complete constraint; try it out:
 - ▶ <https://sat-modulo-symmetries.readthedocs.io>
 - ▶ <https://github.com/markirch/sat-modulo-symmetries>
- a SAT-based exploration of small cases can inspire general results;
- highly symmetric hypergraphs are quite hard for SMS;
- also hard for coloring, in particular when not colorable (extremal);

- [Ber89] Claude Berge.
On the chromatic index of a linear hypergraph and the Chvátal conjecture.
Annals of the New York Academy of Sciences, 555(1):40–44, 1989.
- [Für86] Zoltán Füredi.
The chromatic index of simple hypergraphs.
Graphs and Combinatorics, 2(1):89–92, 1986.
- [Hin81] Neil Hindman.
On a conjecture of Erdős, Faber, and Lovász about n -colorings.
Canadian Journal of Mathematics, 33(3):563–570, 1981.
- [KKK⁺21] Dong Yeap Kang, Tom Kelly, Daniela Kühn, Abhishek Methuku, and Deryk Osthus.
A proof of the Erdős-Faber-Lovász conjecture, 2021.
- [KPS23] Markus Kirchweger, Tomáš Peitl, and Stefan Szeider.
Co-certificate learning with SAT modulo symmetries.

In *Proceedings of the 34th International Joint Conference on Artificial Intelligence, IJCAI 2023*. AAAI Press/IJCAI, 2023.

To appear.

[KS21] Markus Kirchweger and Stefan Szeider.

SAT modulo symmetries for graph generation.

In *27th International Conference on Principles and Practice of Constraint Programming (CP 2021)*, LIPIcs, page 39:1–39:17. Dagstuhl, 2021.

[RAP14] David Romero and Federico Alonso-Pecina.

The Erdős–Faber–Lovász conjecture is true for $n \leq 12$.

Discrete Mathematics, Algorithms and Applications, 06(03):1450039, 2014.